



Phylogenetic patterns of trait and trait plasticity evolution: Insights from amphibian embryos

Rick A. Relyea,^{1,2} Patrick R. Stephens,³ Lisa N. Barrow,⁴ Andrew R. Blaustein,⁵ Paul W. Bradley,⁶ Julia C. Buck,⁷ Ann Chang,⁸ James P. Collins,⁹ Brian Crother,¹⁰ Julia Earl,¹¹ Stephanie S. Gervasi,¹² Jason T. Hoverman,¹³  Oliver Hyman,⁹ Emily Moriarty Lemmon,¹⁴  Thomas M. Luhring,¹⁵ Moses Michelson,¹⁴ Chris Murray,¹⁶ Steven Price,¹⁷ Raymond D. Semlitsch,¹⁸ Andrew Sih,¹⁹ Aaron B. Stoler,¹ Nick VandenBroek,¹⁰ Alexa Warwick,¹⁴ Greta Wengert,¹⁹ and John I. Hammond⁴

¹Department of Biological Sciences, Rensselaer Polytechnic Institute, Darrin Fresh Water Institute, Troy, New York 12180

²E-mail: relyer@rpi.edu

³Odum School of Ecology, University of Georgia, Athens, Georgia 30602

⁴Department of Biology, University of New Mexico, Albuquerque, New Mexico 87131

⁵Department of Integrative Biology, Oregon State University, Corvallis, Oregon 97331

⁶Department of Biology, University of San Diego, San Diego, California 92110

⁷Marine Science Institute, University of California, Santa Barbara, California 93106

⁸Museum of Vertebrate Zoology, University of California, Berkeley, California 94720

⁹School of Life Sciences, Arizona State University, Tempe, Arizona 85287

¹⁰Department of Biological Sciences, Southeastern Louisiana University, Hammond, Louisiana 70402

¹¹Department of Natural Resource Ecology and Management, Oklahoma State University, Stillwater, Oklahoma 74078

¹²Monell Chemical Senses Center, Philadelphia, Pennsylvania 19104

¹³Department of Forestry & Natural Resources, Purdue University, West Lafayette, Indiana 47907

¹⁴Department of Biology, Florida State University, Tallahassee, Florida 32306

¹⁵School of Biological Sciences, University of Nebraska, Lincoln, Nebraska 68588

¹⁶Department of Biology, Tennessee Technological University, Cookeville, Tennessee 38505

¹⁷Department of Forestry, University of Kentucky, Lexington, Kentucky 40546

¹⁸Division of Biological Sciences, University of Missouri, Columbia, Missouri 65211

¹⁹College of Biological Sciences, University of California-Davis, Davis, California 95616

Received February 11, 2017

Accepted December 27, 2017

Environmental variation favors the evolution of phenotypic plasticity. For many species, we understand the costs and benefits of different phenotypes, but we lack a broad understanding of how plastic traits evolve across large clades. Using identical experiments conducted across North America, we examined prey responses to predator cues. We quantified five life-history traits and the magnitude of their plasticity for 23 amphibian species/populations (spanning three families and five genera) when exposed to no cues, crushed-egg cues, and predatory crayfish cues. Embryonic responses varied considerably among species and phylogenetic signal was common among the traits, whereas phylogenetic signal was rare for trait plasticities. Among trait-evolution models, the Ornstein–Uhlenbeck (OU) model provided the best fit or was essentially tied with Brownian motion. Using the best fitting model, evolutionary rates for plasticities were higher than traits for three life-history traits and lower for two.

These data suggest that the evolution of life-history traits in amphibian embryos is more constrained by a species' position in the phylogeny than is the evolution of life history plasticities. The fact that an OU model of trait evolution was often a good fit to patterns of trait variation may indicate adaptive optima for traits and their plasticities.

KEY WORDS: *Anaxyrus*, *Hyla*, *Lithobates*, *Pseudacris*, phylogenetic inertia, *Rana*.

Most organisms in nature experience environmental variation. In response to this variation, many species have evolved traits that are induced by the environment to produce alternative, adaptive phenotypes (i.e., phenotypically plastic traits). As a result, we have an excellent understanding of plasticity in numerous traits from a wide range of plants, animals, fungi, protists, and bacteria and from a variety of biological disciplines including genetics, molecular biology, developmental biology, and ecology (see reviews by Schlichting and Pigliucci 1998; West-Eberhard 2003; DeWitt and Scheiner 2004; Callahan et al. 2008; Murren et al. 2014, 2015). Although adaptive plasticity is ubiquitous, our understanding in any particular system typically comes from intensive investigations on a limited set of species within a clade. What has been missing in this endeavor has been the examination of broad patterns of plasticity evolution across a clade within a phylogenetic perspective.

The study of phylogenetics has enjoyed a rich history in biology as a way of understanding the evolutionary relationships among species (Grant 1986; Harvey and Pagel 1991; Schluter 2000; Berendonk et al. 2003; Felsenstein 2004). There has been increased interest in using phylogenies as a map onto which the species' ecology and traits can be overlaid to determine whether species possess similar traits due to a shared history or due to similar selective forces that produce convergence (Losos 1990; Winemiller 1991; Cadle and Greene 1993; Losos et al. 1998; McPeck and Brown 2000; Price et al. 2000; Webb et al. 2002; Stephens and Wiens 2004, see Special Feature of Ecology, 2012 Supplement).

Given the power of a phylogenetic perspective, it is not surprising that there have been repeated calls for the integration of phylogenetics and phenotypic plasticity beyond the traditional examination of closely related populations or congeners (Doughty 1995; Diggle 2002; Pigliucci et al. 2003; Murren et al. 2015). By assessing the directions and magnitudes of adaptive trait changes for a large number of related species, we can address important evolutionary questions. One question of particular interest is whether, like most morphological and ecological traits, trait plasticity exhibits phylogenetic signal (defined as a pattern where trait disparity scales with the phylogenetic distance that separates species; Blomerg et al. 2003; Revell et al. 2008; Losos 2008). Environmental variation poses diverse challenges to species performance, but it is clear that species can evolve a variety of effective plastic strategies (Boersma et al. 1998; Pigliucci 2001; West-Eberhard 2003). As a result, we might expect different clades

within a phylogenetic group to either evolve particular types of plasticity in response to environmental variation (i.e., morphological, behavioral, or physiology) or evolve a range of unique combinations of nonplastic phenotypes that are similarly effective in the particular environment. The existence of phylogenetic signal can represent a level of phylogenetic inertia that could constrain how traits or trait plasticity can evolve, in the sense that close relatives will be constrained to have similar trait values due to shared descent. Although we know a good deal about the phylogenetic signal of traits (see above citations), we know considerably less about the presence of phylogenetic signal in the plasticity of traits (Pigliucci et al. 1999; Pollard et al. 2001; Pigliucci et al. 2003; Thaler and Karban 1997; Colbourne et al. 1997; Gomez-Mestre et al. 2008; Burns and Strauss 2012).

If we could map traits and trait plasticity onto a phylogeny, we could also examine questions about rates of evolution. We could compare rates of evolution among different types of traits (e.g., behavior, morphology, life history) and different magnitudes of trait plasticity. Rates of evolution have been investigated in many different studies of species' traits, but there appear to be no studies that have compared the rates of trait evolution to rates of plasticity evolution. Thus, it remains an open question whether traits evolve slower or faster than trait plasticity. Answering this question should provide insights into the role that phylogenetic relationships play in constraining the evolution of phenotypically plastic traits, and whether species traits or trait plasticity will be able to shift more quickly in the face of environmental change.

In this study, we examined the phylogenetic patterns of phenotypic plasticity using amphibians, a model system that has become well known for exhibiting predator-induced behavioral, morphological, and life-history traits (Van Buskirk 2002; Miner et al. 2005; Relyea 2007). Our focus was to examine how amphibian embryos respond to predators. Predators of eggs (e.g., crayfish, snakes, and leeches) induce several species of amphibians to hatch earlier and with a smaller mass or at a less-developed stage, whereas other species are unresponsive (Warkentin 1995; Chivers et al. 2001; Laurila et al. 2001, 2002; Johnson et al. 2003; Saenz et al. 2003; Orizaola and Braña 2004; Vonesh 2005; Gomez-Mestre et al. 2008; Anderson and Brown 2009; Segev et al. 2015). From the studies that have been conducted on amphibian embryos (encompassing 12 families), one can conclude that predator-induced developmental plasticity exists in some, but not all species. Because most of these experimental efforts thus far have examined only one or two species at a time (using a

variety of experimental conditions and predator species), we have relatively few comparable data on the directions and magnitudes of these responses. We also lack a general understanding of the phylogenetic constraints and ecological conditions under which these responses have evolved (but see Gomez-Mestre et al. 2008). Clearly, a phylogenetic approach is well suited to examine this question.

We addressed the following hypotheses: (1) Amphibian embryos will respond to predator cues by hatching earlier, less developed, and at a smaller size, (2) different species of amphibian embryos will differ in their traits and trait plasticity in response to predation risk, (3) there will be phylogenetic signal in the traits and trait plasticity of amphibian embryos, and (4) the rates of evolution will be similar between traits and trait plasticity, when estimated using the best fitting model of character evolution.

Methods

THE CHALLENGES

When taking a phylogenetic approach to quantifying ecologically important traits, a number of challenges arise (for both plastic and nonplastic traits; i.e., constitutive traits). The first challenge is deciding upon the rearing conditions. Many biologists would prefer to quantify traits as they appear in nature, yet this method would confound species-level variation with environmental factors, including differences in temperature, pH, dissolved oxygen, and resources. Thus, investigators doing comparative work on plastic traits have assessed species under common-garden conditions to ensure that observed phenotypic differences can be attributed to genetic differences among species (Pigliucci et al. 1999, 2003; Colbourne 1997; Thaler and Karban 1997; Richardson 2001a,b, 2002a,b). One limitation is that species living under common-garden conditions might not exhibit the same magnitude of plasticity that they exhibit in nature. Given the extreme logistical difficulties of raising a large number of species under a wide range of environmental conditions, the common-garden approach is a necessary compromise that arises from the need to balance between these challenges. However, because temperature is one environmental condition of particular concern, we assessed the potential bias of temperature on plastic responses by rearing three of the species under two different temperatures.

The second challenge in taking a phylogenetic approach is to select populations that best represent a species. For most species, there is population-level variation in traits, especially in cosmopolitan species. However, comparative studies generally make the assumption that interspecific variation is larger than intraspecific variation. Because of the substantial challenge of assessing a large number of species, it is not feasible to sample multiple populations throughout each species' range (although

this is an interesting question that could be examined in future studies). To circumvent this problem, we attempted to minimize nonrepresentative phenotypes by collecting species in their most common type of habitat and avoiding the collection of individuals from the extremes of a species' range. However, we also assessed the impact of intraspecific variation on our conclusions by quantifying the plasticity of two distant populations from each of three cosmopolitan species (*Lithobates catesbeianus*, *Anaxyrus americanus*, and *Hyla versicolor*).

THE EXPERIMENTS

All animals were collected as newly oviposited egg masses by the different laboratory groups from around the United States (Table SA1). Once collected, eggs were transported back to the local laboratory and held at 21°C on a 12-h light:12-h dark cycle. Because the experiments were conducted in several laboratories around the United States, all laboratories used identical water mixtures (i.e., 20 L of deionized water mixed with 75 g of NaCl, 2.1 g of MgSO₄, 1.05 g of KCl, 4.2 g of NaHCO₃, and 2.8 g of CaCl).

Each egg-hatching experiment was conducted using a completely randomized design with three treatments and six replicates of each treatment for a total of 18 experimental units. The experimental units were Petri dishes (plastic 100 × 20 mm) containing 50 mL of water. To determine whether the embryos exhibited plastic responses to the three environments, we waited until the eggs approximately reached gastrulation stage (Gosner stage 12; Gosner 1960) and separated 190 eggs from each collection of clutches. In all experiments, we randomly assigned 10 eggs to each of the 18 Petri dishes and preserved an additional 10 eggs in buffered 10% formalin to confirm the developmental stage of the eggs at the start of each experiment. In one case (the Oregon population of *Anaxyrus boreas*), the eggs were collected at a slightly later stage (Gosner stage 15).

Our three treatments were control (i.e., no predator cues), crushed conspecific eggs, and crayfish-consumed conspecific eggs. The cues for each environment were generated using 1 L of water that had either no predator cues (i.e., control), eggs that were crushed by hand, or eggs that were consumed by a red swamp crayfish (*Procambarus clarkii*). We chose a crayfish because as a group they are common consumers of amphibian eggs. We chose the red swamp crayfish in particular because it has a large native range in North America and an even larger introduced range (<https://nas.er.usgs.gov/viewer/omap.aspx?SpeciesID=217>). The implicit assumption is that the antipredator responses induced by chemical cues of this crayfish would be similar to those induced by other crayfish that coexist with the various amphibian species. Unfortunately, there is probably no single egg predator species that coexists with every anuran species in North America.

For the crushed and consumed-egg treatments, we used three, five, or seven eggs to generate the cues, depending on egg size; the smaller the eggs of a given species, the more eggs were crushed or consumed to provide all species with an approximately equal amount of egg biomass that could produce chemical cues. This is important because the mass of prey fed to a predator affects the strength of the prey's response (Schoeppner and Relyea 2008). To increase the likelihood that we would have a predator consume all eggs within 30 min, we set up three, 1-L crayfish containers; the container with the highest predation (i.e., usually consuming all eggs) was used for the cue source. All cues were added 30 min after being generated in the crushed and crayfish treatments. For both treatments, we removed a sample of the water containing the cues and avoided picking up any organic matter (e.g., pieces of destroyed eggs or feces from the crayfish). When adding the cues to the dishes, we removed 25 mL of water from each dish every 12 h and replaced it with 25 mL of new cue water. After the cues were added to the dishes, we added new water to the cue-generating containers to return their volume to 1 L.

As the embryos approached hatching, we checked the dishes every 4 h to determine the time of hatching (i.e., the time at which an individual successfully leaves its egg). As each animal hatched, we recorded the time to hatching and then euthanized and preserved the individual in buffered 10% formalin to later determine the Gosner stage and mass at hatching. Across all experiments, embryo survival ranged from 58 to 97% (median = 89%). At the end of the experiment, preserved hatchlings from all laboratories were shipped to the University of Pittsburgh, where we quantified the mean Gosner stage at the start of each embryonic experiment, mean developmental stage at the time of hatching, and mean individual mass. Using these response variables, we calculated developmental rate (i.e., [stage at hatching – initial stage] ÷ time to hatching) and growth rate (i.e., mass at hatching ÷ time to hatching).

ADDITIONAL EXPERIMENTS MANIPULATING TEMPERATURE

To assess the impact of different rearing temperatures on our conclusions, we examined the effects of different temperatures on the traits and trait plasticity of the embryos. To do this, we tested three of the species (one species from each family: *A. americanus*, *H. versicolor*, and *L. clamitans*) in identical experiments as described above, but at a second temperature (19°C).

STATISTICAL METHODS FOR ASSESSING HOW EACH SPECIES RESPONDED TO THE ENVIRONMENT

We assessed the plasticity of five traits for each species: mass, growth rate, time to hatching, stage at hatching, and developmental rate. We began by taking the mean of all individuals in an experimental unit for each response variable. In general, the mean

time to hatching, growth rate, and developmental rate were all normally distributed; only the *Anaxyrus fowleri* was nonnormally distributed, but nonparametric tests gave similar results as parametric tests. Because analysis of variance is robust to violations of this assumption, we used a multivariate analysis of variance (MANOVA) for each species followed by subsequent univariate tests for each response variable. Stage at hatching was generally nonnormally distributed; moreover, eight of the 23 species exhibited no variation in this trait, so this variable was removed from the MANOVA for these species (Table SA3). All other data were rank-transformed. To test for trait induction due to an exposure to crushed or consumed eggs, we used planned contrasts between the control and crayfish treatments and between the control and crushed conspecific treatments.

To determine if temperature interacted with plasticity for the three species that we raised at two temperatures, we used a MANOVA to test for effects of temperature, treatment, and their interaction. Response variables were generally normal within temperatures. All three species exhibited significant multivariate effects of temperature (Table SA4) and one of the species exhibited a multivariate temperature-by-treatment interaction. However, this interaction was driven by a difference between responses to crayfish versus crushed eggs at the two different temperatures. Our interest was in examining plasticity in control versus crayfish and control versus crushed eggs and these specific plastic responses did not interact with temperature (all $P > 0.19$) for any of the traits in any of the three species. Because the magnitude of the plastic responses did not change as a function of the two temperatures, we used the data from the 21°C experiment to match all of the other species. Although such results are encouraging, we cannot assess whether temperature might cause interactive effects on the magnitude of plasticity with the other species.

STATISTICAL METHODS FOR QUANTIFYING TRAIT PLASTICITY

To address our questions about phylogenetic patterns of plasticity, we first had to determine how to quantify plasticity and how to interpret the outcomes when plasticity is quantified in different ways. We included measures of plasticity as an absolute measure (e.g., trait mean for the control treatment – trait mean for the crayfish treatment), plasticity that is proportional to the grand mean (e.g., [trait mean for the control treatment – trait mean for the crayfish treatment] ÷ grand mean), and plasticity that is proportional to the pooled SD, using the meta-analytic Hedge's G (e.g., [trait mean for the control treatment – trait mean for the crayfish treatment] ÷ pooled SD; Hedges 1981). These plasticity measures included both direction and magnitude. For the pooled SDs, we defined SD_1 as the pooled SD for the control and crushed environments and SD_2 as the pooled SD for the control and crayfish environments.

Given that some of our plasticity measures used the grand mean or a pooled SD as a denominator, we also decided it was important to determine whether these denominators contained any phylogenetic signal that would cause a spurious result when we looked for a phylogenetic signal of plasticity. As a result, we assessed phylogenetic signal in the grand mean and in the pool SDs. In this way, when we assessed phylogenetic signal using plasticity metrics that included grand means and pooled SDs, we could assess the contribution of signal from both the numerator and denominator.

PHYLOGENETIC COMPARATIVE ANALYSES

Species phylogenetic relationships were obtained from a published amphibian phylogeny (Pyron and Wiens 2013), pruned down to include only the species found in our study. Analyses with this tree were conducted using values averaged across populations to produce a species-level estimate when multiple populations were measured. Phylogenetic signal was quantified using Blomberg's K (Blomberg et al. 2003). The statistical significance of phylogenetic signal was assessed using a randomization test described by Blomberg and Garland (2002) implemented in the R library *picante* (Kembel et al. 2010). Tests were performed using 10,000 randomizations. In all cases, we consider P values significant at values of ≤ 0.05 . Our measurements of developmental stage included a few species that expressed zero plasticity. We calculated rate and signal for developmental stage plasticity both including and excluding species that exhibited zero plasticity and the results were quantitatively similar and qualitatively identical in both sets of analyses. As a result, we report results using all species.

For each trait and plasticity measure, we compared the fits (i.e., AICc scores) of three models of continuous character evolution: (1) the Brownian motion (BM) model of character evolution, which is based on models of random particle diffusion in a liquid, which is a widely used neutral model of character evolution under genetic drift and the model implicitly assumed by many phylogenetic comparative methods; (2) the Ornstein–Uhlenbeck (OU) model of character evolution (Hansen 1997, Butler & King 2004), which models character evolution as a random walk with a central tendency (i.e., a stabilizing force); and (3) the early burst (EB) model that models character evolution as a random walk, the rate of which decreases as you move from the root of the tree to the tips. Evolutionary rate was measured using sigma squared estimated from models fitted using the R package *Geiger* version 2.0.6 (Harmon et al. 2008; Pennell et al. 2014). For OU models, we estimated α (i.e., the strength of attraction toward a central tendency) to evaluate the degree to which OU models differed from BM. We also estimated the phylogenetic half-life, which is a measure of how fast species would be expected to approach the evolutionary optimum.

The three models were first fitted using raw trait data and measures of plasticity. However, this produced rate estimates that were highly correlated with the order of magnitude of the range of trait variation (Fig. SA2). For example, if one trait varied between 1 and 10 and another varied between 10 and 100, the latter trait would show a much higher rate estimate even though in proportion to their size both traits are likely evolving at similar rates. The typical method of dealing with this issue would be to estimate rates for log-transformed traits (Ackerly 2009). However, most of our measures of plasticity contained negative values for some species, which cannot be log transformed. Transforming the absolute values of trait plasticity was considered. However, this would have discarded information on the direction of plasticity, and would have greatly inflated rate estimates for trait plasticity that happened to show positive or negative values close to zero. Instead, we divided all trait and trait plasticity values by the minimum value that any species exhibited, and used these ratio-transformed trait and trait plasticity values to estimate evolutionary rate. Like log-transformed traits, this yielded trait ranges that were wide when the maximum species trait values were multiples of minimum trait values, but that also preserved information on directionality. Only the results using ratio-transformed trait and plasticity values are reported here.

Results

TRAIT PLASTICITY IN EACH SPECIES

Our first analysis conducted MANOVAs on the five life-history traits for each of the 23 experiments (Table SA2; Figs. 1 and 2). Of the 23 experiments, nine had significant multivariate tests and one had a nearly significant multivariate test (i.e., $P < 0.07$).

We then examined the patterns of responses to crayfish consuming eggs versus the control, which should represent the highest risk environment. For time to hatching, four species hatched earlier with crayfish ($P < 0.05$), six species hatched later ($P < 0.05$), and the remaining 13 species exhibited no significant response. For mass at hatching, three species hatched at a smaller mass with crayfish, one species had a greater mass, and 20 species exhibited no response. For growth rate, three species grew at a slower rate with crayfish, while the other 20 species exhibited no response. For stage at hatching, three species were less developed with crayfish, two species were more developed, and 18 species exhibited no response (including the eight species that exhibited zero variation in this trait). For developmental rate, five species had a slower developmental rate with crayfish, four species had a faster developmental rate, and 14 species exhibited no response.

Next, we examined the patterns of responses to crushed eggs versus the control (Table SA2; Figs. 1 and 2). For time to hatching, two species hatched earlier with crushed eggs, two species hatched




Figure 1. Time to hatching, mass at hatching, and Gosner stage at hatching for 20 species of amphibian embryos when exposed to cues from either no predator, crushed conspecific eggs, or crayfish consuming conspecific eggs. To test for population effects, three species were selected from two locations. Abbreviations refer to the states in which the animals were collected. Asterisks indicate species in which crushed-egg cues differed from the control; crosses indicate species in which crayfish cues differed from the control ($P < 0.05$). Data are means \pm 1 SE.

later, and the remaining 19 species exhibited no response. For mass at hatching, three species hatched at a smaller mass with crushed eggs, one species had a greater mass, and 19 species exhibited no response. For growth rate, three species grew at a slower rate with crushed eggs, whereas the other 20 species exhibited no response. For stage at hatching, two species were less developed with crushed eggs, three species were more developed, and 18 species exhibited no response (including the eight species that exhibited zero variation in this trait). For developmental rate, two species had a slower developmental rate with crushed eggs,

three species had a faster developmental rate, and 18 species exhibited no response.

THERMAL SENSITIVITY IN PLASTICITY AMONG THREE FOCAL SPECIES

For three of the species, we raised the embryos at two different temperatures to determine the importance of temperature in altering the traits and the plasticity of the traits (i.e., as determined by a treatment-by-temperature interaction; Table A3). For *A. americanus* and *L. clamitans*, temperature had a




Figure 2. Growth rate (i.e., mass ÷ time to hatch) and developmental rate (i.e., Gosner stage ÷ time to hatch) for 20 species of amphibian embryos when exposed to cues from either no predator, crushed conspecific eggs, or crayfish consuming conspecific eggs. To test for population effects, three species were selected from two locations. Abbreviations refer to the states in which the animals were collected. Asterisks indicate species in which crushed-egg cues differed from the control; crosses indicate species in which crayfish cues differed from the control ($P < 0.05$). Data are means \pm 1 SE.

multivariate effect that was driven by four of the five embryo traits, but temperature did not alter the magnitudes of plasticity. For *H. versicolor*, however, there were multivariate effects of temperature, treatment, and their interaction. The interaction was driven by the mass at hatching and stage at hatching. However, subsequent analyses indicated that contrasts of interest (i.e., the magnitude of plasticity for the control vs. crushed-egg treatments and the magnitude of plasticity for the control vs. consumed-egg treatments) did not differ with temperature.

PHYLOGENETIC SIGNAL IN TRAITS

Our analysis of phylogenetic signal began by testing for phylogenetic signal in the five embryonic traits within each of the

three environments (control, crayfish cues, and crushed-egg cues; Tables 1–5, SA4–8). Time to hatching did not exhibit statistically significant phylogenetic signal in any environment. Mass at hatching, stage at hatching, and growth rate all exhibited significant phylogenetic signal in all three environments. Development rate exhibited significant phylogenetic signal in two environments and nearly significant ($P = 0.054$) in the third. Across all 15 analyses of phylogenetic signal in embryonic traits, phylogenetic signal was significant in 73% of the tests (Table 6).

PHYLOGENETIC SIGNAL IN TRAIT PLASTICITY

We next tested for phylogenetic signal in the plasticity of the five embryonic traits using multiple measures of plasticity

Table 1. Tests of phylogenetic signal and evolutionary rates for time to hatching of amphibian embryos when raised in three inducing environments.

	Blomberg's <i>K</i>	<i>P</i> -value	BM AICc	OU AICc	EB AICc
(A) Trait					
Control	0.339	0.076	49.4	45.1	52.2
Crushed-egg cues	0.338	0.082	50.2	45.6	53.1
Crayfish cues	0.313	0.116	54.0	48.6	56.9
(B) Trait plasticity					
(Control – Crushed)	0.189	0.417	49.3	35.4	52.2
(Control – Crayfish)	0.131	0.700	49.9	30.0	52.8
(Control – Crushed) ÷ Grand mean	0.186	0.421	49.4	35.3	52.2
(Control – Crayfish) ÷ Grand mean	0.170	0.488	49.5	34.2	52.4
(Control – Crushed) ÷ Pooled SD ₁	0.197	0.393	43.7	30.2	46.5
(Control – Crayfish) ÷ Pooled SD ₂	0.135	0.676	61.0	41.6	63.8
(C) Grand mean and pooled SD					
Grand mean	0.331	0.092	51.0	46.3	53.9
Pooled SD ₁	0.186	0.437	86.1	73.1	89.0
Pooled SD ₂	0.261	0.215	77.1	70.2	80.0

The analyses tested (A) traits and (B) trait plasticity of the embryos. To better understand how phylogenetic signal was affected by the numerators and denominators in the plasticity estimates, we also assessed phylogenetic signal in the (C) grand mean and pooled SDs. Phylogenetic signal was tested using Blomberg's *K* and its associated *P*-value (bold font indicates significant tests; $P < 0.05$). Three models of continuous character evolution were used to determine which model produced the best fit to the data (based on AICc scores): Brownian motion (BM), Ornstein–Uhlenbeck (OU), and early burst (EB). Models with the lowest AICc values are shown in bold font.

(Tables 1–5, SA4–8). For time to hatching and developmental rate, none of the six tests exhibited significant phylogenetic signal. For mass at hatching and growth rate, two of the six measures of plasticity (Control – Crushed and Control – Crayfish) exhibited significant signal. For stage at hatching, only one of the six measures ([Control – Crushed]/SD₁) exhibited significant signal. Across all 30 analyses of phylogenetic signal in trait plasticity, phylogenetic signal was significant in 17% of the tests (Table 6).

MODELS OF TRAIT EVOLUTION AND EVOLUTIONARY RATES OF TRAITS VERSUS TRAIT PLASTICITY

For time to hatching and developmental rate, the OU model was a better fit for both traits and their plasticity. For the other three traits, the OU model was essentially tied with BM, with comparisons of the two models exhibiting delta AICc scores of less than two. It is notable that the best fitting model of trait evolution was always the same for both trait means and plasticity measures derived from them for any given life-history trait. Values of α estimated for OU models in cases where the OU model was a better fit were higher than those estimated in cases where BM was a better fit (Table 7). For time to hatching and developmental rate, plasticity tended to show a higher evolutionary rate than trait means under the best fitting (i.e., OU) model of character evolution. For stage at hatching, plasticity showed higher evolutionary rates than trait means regardless of whether sigma squared was estimated using a BM or OU model. For mass at hatching and growth rate,

the opposite held true; plasticity tended to show a lower evolutionary rate than trait means. We note that the evolutionary rates for the pooled SD generally showed a greatly inflated evolutionary rate, but the method employed to measure evolutionary rates was designed for species trait means, not SDs.

Discussion

In documenting the life-history responses to predator cues in three families of amphibians, we discovered that responses varied a great deal among species. Moreover, we found that the traits frequently exhibited phylogenetic signal, whereas trait plasticity rarely exhibited phylogenetic signal. Finally, we discovered that evolutionary rates were more rapid for traits than trait plasticity for two life-history traits, but less rapid for three life-history traits. Next, we expand on each of these discoveries.

THE PHENOTYPIC RESPONSES OF EMBRYOS COMPARED TO THE LITERATURE

More than 30 embryonic plasticity studies have been conducted with amphibians over the past two decades to quantify their life-history responses to predators under a variety of experimental conditions (Segev et al. 2015). These studies have found a range of predator-induced responses that suggest that different species have evolved different magnitudes of embryonic responses to egg predators, damaged eggs, and larval predators (including a

Table 2. Tests of phylogenetic signal and evolutionary rates for mass at hatching of amphibian embryos when raised in three inducing environments.

	Blomberg's <i>K</i>	<i>P</i> -value	BM AICc	OU AICc	EB AICc
(A) Trait					
Control	0.737	0.003	94.6	96.6	97.4
Crushed-egg cues	0.730	0.004	92.6	94.5	95.4
Crayfish cues	0.733	0.003	96.2	98.0	99.0
(B) Trait plasticity					
(Control - Crushed)	0.473	0.028	72.5	71.8	75.4
(Control - Crayfish)	0.485	0.037	93.2	93.3	96.1
(Control - Crushed) ÷ Grand mean	0.160	0.566	44.7	28.2	47.6
(Control - Crayfish) ÷ Grand mean	0.179	0.428	77.3	63.5	80.1
(Control - Crushed) ÷ Pooled SD ₁	0.221	0.288	65.9	55.4	68.8
(Control - Crayfish) ÷ Pooled SD ₂	0.228	0.251	62.3	53.1	65.2
(C) Grand mean and pooled SD					
Grand mean	0.735	0.003	94.4	96.3	97.2
Pooled SD ₁	0.605	0.008	131.1	132.3	134.0
Pooled SD ₂	0.501	0.058	150.4	150.6	153.3

The analyses tested (A) traits and (B) trait plasticity of the embryos. To better understand how phylogenetic signal was affected by the numerators and denominators in the plasticity estimates, we also assessed phylogenetic signal in the (C) grand mean and pooled SDs. Phylogenetic signal was tested using Blomberg's *K* and its associated *P*-value (bold font indicates significant tests; $P < 0.05$). Three models of continuous character evolution were used to determine which model produced the best fit to the data (based on AICc scores): Brownian motion (BM), Ornstein-Uhlenbeck (OU), and early burst (EB). Models with the lowest AICc values are shown in bold font.

complete lack of response). However, a major challenge in interpreting the patterns of induced responses among species is that nearly all of these studies have examined one species at a time, although a few have examined two species. The power of the current study's approach is that all 20 species were raised under identical conditions with the same suite of environmental manipulations, making it possible to attribute differences in response to the species and not differences in experimental design (see also Gomez-Mestre et al. 2008). However, as noted earlier, an important constraint in doing large comparisons among species raised under identical conditions is that it becomes difficult or impossible to know whether the fundamental conclusions would change under different abiotic conditions including differences in temperature, pH, and per capita food rations. This constraint also applies to comparing all past studies that have raised one or more species under controlled conditions, whether in the laboratory or in outdoor venues.

In regard to the time it takes for an embryo to hatch, one would predict an adaptive response to egg predators would be for embryos to hatch sooner and avoid the predator, although this would likely come at the cost of reduced growth and development of the newly hatched animal. Consistent with this prediction, a number of studies have observed earlier hatching times (e.g., Johnson et al. 2003; Touchon et al. 2006; Segev et al. 2015), but a substantial number of studies have not (e.g., Schalk et al. 2002; Anderson and Petranka 2003; Dibble et al. 2009). Similarly, even when cues

from egg predators induce earlier hatching, the expected trade-off of smaller, less developed hatchlings sometimes is observed (Johnson et al. 2003; Gomez-Mestre et al. 2008), but many times the response variables are simply not measured. In our experiments, however, there was no evidence of a consistent trade-off; several species that were induced to exhibit significantly shorter or longer times to hatching did not exhibit significant changes in their mass at hatching or stage at hatching. Of course, a number of constraints could prevent such a response from evolving.

In our set of 23 experiments, embryos exposed to crayfish induced earlier time to hatching in four experiments, a later time to hatching in six experiments, and no change in 13 experiments. For comparison, there have been three studies of amphibian embryos raised with and without crayfish cues and all of them have observed earlier times to hatching with crayfish cues (using *L. sphenoccephala* and *L. clamitans*; Johnson et al. 2003; Saenz et al. 2003; Anderson & Brown 2009). In our study, *L. clamitans* also tended to hatch earlier as well, but the difference was not significantly different from the control ($P = 0.19$). We have no way of assessing potential publication bias in which studies may have observed no response to crayfish cues and not published the results. Of course, one would not expect every species to respond to a given predator in the same way (Relyea 2007) and plastic responses may depend on a wide variety of factors including abiotic conditions, ontogeny, and whether the predators consumes both eggs and larvae of amphibians (as crayfish can do). As noted

Table 3. Tests of phylogenetic signal and evolutionary rates for developmental stage to hatching of amphibian embryos when raised in three inducing environments.

	Blomberg's <i>K</i>	<i>P</i> -value	BM AICc	OU AICc	EB AICc
(A) Trait					
Control	1.079	0.000	-32.1	-30.4	-29.3
Crushed-egg cues	1.167	0.000	-33.2	-31.2	-30.3
Crayfish cues	1.131	0.001	-32.4	-30.6	-29.6
(B) Trait plasticity					
(Control – Crushed)	0.173	0.547	52.9	38.6	55.8
(Control – Crayfish)	0.260	0.473	67.5	60.9	70.3
(Control – Crushed) ÷ Grand mean	0.197	0.492	56.5	44.7	59.4
(Control – Crayfish) ÷ Grand mean	0.251	0.483	66.3	59.1	69.2
(Control – Crushed) ÷ Pooled SD ₁	0.479	0.014	12.9	13.5	15.8
(Control – Crayfish) ÷ Pooled SD ₂	0.234	0.333	56.3	47.7	59.2
C. Grand mean and pooled SD					
Grand mean	1.135	0.000	-32.8	-30.9	-30.0
Pooled SD ₁	0.226	0.279	32.1	22.2	35.0
Pooled SD ₂	0.317	0.086	NE	NE	NE

The analyses tested (A) traits and (B) trait plasticity of the embryos. To better understand how phylogenetic signal was affected by the numerators and denominators in the plasticity estimates, we also assessed phylogenetic signal in the (C) grand mean and pooled SDs. Phylogenetic signal was tested using Blomberg's *K* and its associated *P*-value (bold font indicates significant tests; $P < 0.05$). Three models of continuous character evolution were used to determine which model produced the best fit to the data (based on AICc scores): Brownian motion (BM), Ornstein–Uhlenbeck (OU), and early burst (EB). Models with the lowest AICc values are shown in bold font (NE, not estimable due to a denominator of zero).

Table 4. Tests of phylogenetic signal and evolutionary rates for growth rate of amphibian embryos when raised in three inducing environments.

	Blomberg's <i>K</i>	<i>P</i> -value	BM AICc	OU AICc	EB AICc
(A) Trait					
Control	0.5948	0.0069	94.2	93.9	97.0
Crushed-egg cues	0.6049	0.0069	91.4	91.1	94.3
Crayfish cues	0.5863	0.0079	94.3	93.6	97.1
(B) Trait plasticity					
(Control – Crushed)	0.6188	0.0061	46.9	49.2	49.7
(Control – Crayfish)	0.4755	0.0306	60.8	61.3	63.6
(Control – Crushed) ÷ Grand mean	0.2175	0.3597	33.0	22.0	35.8
(Control – Crayfish) ÷ Grand mean	0.2106	0.3055	60.5	49.5	63.4
(Control – Crushed) ÷ Pooled SD ₁	0.2355	0.2500	59.7	50.7	62.5
(Control – Crayfish) ÷ Pooled SD ₂	0.3230	0.0643	52.7	48.6	55.5
(C) Grand mean and pooled SD					
Grand mean	0.5961	0.0075	93.2	92.8	96.0
Pooled SD ₁	0.5325	0.0304	133.7	134.0	136.6
Pooled SD ₂	0.4715	0.0506	129.8	129.0	132.6

The analyses tested (A) traits and (B) trait plasticity of the embryos. To better understand how phylogenetic signal was affected by the numerators and denominators in the plasticity estimates, we also assessed phylogenetic signal in the (C) grand mean and pooled SDs. Phylogenetic signal was tested using Blomberg's *K* and its associated *P*-value (bold font indicates significant tests; $P < 0.05$). Three models of continuous character evolution were used to determine which model produced the best fit to the data (based on AICc scores): Brownian motion (BM), Ornstein–Uhlenbeck (OU), and early burst (EB). Models with the lowest AICc values are shown in bold font.

Table 5. Tests of phylogenetic signal and evolutionary rates for developmental rate of amphibian embryos when raised in three inducing environments.

	Blomberg's <i>K</i>	<i>P</i> -value	BM AICc	OU AICc	EB AICc
(A) Trait					
Control	0.353	0.054	61.6	57.7	64.5
Crushed-egg cues	0.422	0.017	56.1	54.1	58.9
Crayfish cues	0.374	0.037	57.9	54.7	60.7
(B) Trait plasticity					
(Control – Crushed)	0.076	0.954	71.6	41.2	74.4
(Control – Crayfish)	0.131	0.713	62.9	42.7	65.8
(Control – Crushed) ÷ Grand mean	0.178	0.490	38.3	24.1	41.2
(Control – Crayfish) ÷ Grand mean	0.191	0.386	52.9	40.1	55.8
(Control – Crushed) ÷ Pooled SD ₁	0.193	0.399	42.3	29.6	45.1
(Control – Crayfish) ÷ Pooled SD ₂	0.113	0.818	65.1	42.5	68.0
(C) Grand mean and pooled SD					
Grand mean	0.384	0.029	57.2	54.3	60.1
Pooled SD ₁	0.190	0.388	94.1	80.8	97.0
Pooled SD ₂	0.312	0.089	81.9	76.6	84.7

The analyses tested (A) traits and (B) trait plasticity of the embryos. To better understand how phylogenetic signal was affected by the numerators and denominators in the plasticity estimates, we also assessed phylogenetic signal in the (C) grand mean and pooled SDs. Phylogenetic signal was tested using Blomberg's *K* and its associated *P*-value (bold font indicates significant tests; $P < 0.05$). Three models of continuous character evolution were used to determine which model produced the best fit to the data (based on AICc scores): Brownian motion (BM), Ornstein–Uhlenbeck (OU), and early burst (EB). Models with the lowest AICc values are shown in bold font.

Table 6. A summary of the analyses for significant phylogenetic signal (Blomberg's *K*) in the (A) traits and (B) trait plasticity of amphibian embryos exposed to predatory cues ($\checkmark = P < 0.05$).

	Time to hatching	Mass at hatching	Stage at hatching	Growth rate	Development rate
(A) Trait					
Control		✓	✓	✓	
Crush		✓	✓	✓	✓
Crayfish		✓	✓	✓	✓
(B) Trait plasticity					
Control – Crushed		✓		✓	
Control – Crayfish		✓		✓	
(Control – Crushed) ÷ GM					
(Control – Crayfish) ÷ GM					
(Control – Crushed) ÷ SD ₁			✓		
(Control – Crayfish) ÷ SD ₂					
(C) Grand mean and pooled SDs					
GM		✓	✓	✓	✓
SD ₁ (control, crushed)		✓		✓	
SD ₂ (control, crayfish)					

Detailed statistics can be found in Tables 1–5. To better understand how phylogenetic signal was affected by the numerators and denominators of several of the plasticity estimates, we also assessed phylogenetic signal in the (C) grand means (GM) and pooled SDs (SD₁ and SD₂).

earlier, one of the challenges in exposing a large number of species from across an entire continent to cues from a single predator species is that not every species will coexist with the predator (although it may coexist with close relatives of the predator). As a result, an important caveat to the results of the crayfish cue treat-

ment is that we may have some cases of species not responding to the cue because that do not coexist with the species of crayfish that we used. One implication of this is that the more reliable data may be how the embryos respond to the cues of crushed eggs.

Table 7. Rate parameters estimated for Ornstein–Uhlenbeck (OU) models of character evolution.

	Time to hatching		Mass at hatching		Developmental stage		Growth rate		Developmental rate	
	α	PHL	α	PHL	α	PHL	α	PHL	α	PHL
(A) Trait										
Control	0.038402	18.04998	0.008496	81.58278	0.008088	85.69541	0.015307	45.28156	0.030426	22.7814
Crushed-egg cues	0.038994	17.77575	0.008725	79.43983	0.006898	100.4809	0.015495	44.73403	0.023369	29.66099
Crayfish cues	0.044294	15.64887	0.008853	78.29554	0.007496	92.47209	0.016042	43.20755	0.027609	25.10627
(B) Trait plasticity										
(Control – Crushed)	0.132396	5.235418	0.020449	33.89608	2.713433	0.25545	0.007647	90.64628	2.718282	0.254995
(Control – Crayfish)	0.460493	1.50523	0.019438	35.65952	0.057431	12.06922	0.01821	38.06354	0.24012	2.886666
(Control – Crushed) ÷ Grand mean	0.150743	4.598212	0.183382	3.77979	1.11786	0.620066	0.112611	6.155223	2.718282	0.254995
(Control – Crayfish) ÷ Grand mean	0.195834	3.539464	0.264430	2.621291	0.689952	1.004631	0.140089	4.947922	0.239441	2.894854
(Control – Crushed) ÷ Pooled SD ₁	0.111275	6.229152	0.142578	4.861535	0.01573	44.06521	0.120524	5.751136	2.718282	0.254995
(Control – Crayfish) ÷ Pooled SD ₂	0.316571	2.189548	0.182090	3.806622	1.293636	0.535813	0.040656	17.04901	0.740391	0.936191
(C) Grand mean and pooled SD										
Grand mean	0.039922	17.36265	0.008636	80.25789	0.007356	94.23198	0.015592	44.45448	0.026646	26.01362
Pooled SD ₁	2.718282	0.254995	0.012946	53.54256	0.220752	3.139933	0.01567	44.23534	0.15818	4.382019
Pooled SD ₂	0.081241	8.531972	0.017354	39.94223	NA	NA	0.020833	33.27169	0.043694	15.86359

Alpha (α) is the attraction strength of the evolutionary optimum, phylogenetic half-life (PHL) is a measure of how fast species would be expected to approach the evolutionary optimum. Values of α in bold indicate instances where the OU model had the lowest AICc score of the three models that were considered.

We also observed that embryos exposed to crushed eggs induced earlier time to hatching in two experiments, a later time to hatching in two experiments, and no change in 19 experiments. We only found two studies that have examined amphibian embryo responses to damaged eggs; one study observed the induction of earlier hatching (*Chiromantis hanseanae*; Poo & Bickford 2014), whereas the other study observed no effect (*L. temporaria*; Mandrillon & Saglio 2007). Collectively, these results suggest that amphibian embryos often respond to cues from crayfish eating conspecific eggs and occasionally respond to cues from damaged conspecific eggs.

PHYLOGENETIC SIGNAL IN TRAITS AND TRAIT PLASTICITY

We observed a striking consistency in the occurrence of phylogenetic signal in the life-history traits. Four of the five sets of life-history trait means showed significant signal, and even time to hatching showed levels of signal that could be described as “nearly significant” with P values of less than 0.1 for two of the three trait means. Detecting phylogenetic signal in the life-history traits of a group of species is not particularly surprising. Such patterns have been detected in a wide range of taxa including arboreal-nesting amphibians (Gomez-Mestre et al. 2008), lizards (Brandt and Navas 2011), carabid beetles (Sota & Ishikawa 2004), and plants (Moles et al. 2005).

The more interesting question for the current study is whether there is phylogenetic signal in trait plasticity. One of the most striking discoveries was how rarely trait plasticity exhibited phylogenetic signal across the five life-history variables. From our 30 assessments of phylogenetic signal in life-history traits, only 17% exhibited significant phylogenetic signal. Phylogenetic signal was completely lacking for time to hatching, whereas it was only present in 17–33% of the six comparisons within each type of trait. To evaluate the robustness of our results, we used three metrics: (1) the difference in phenotypic value expressed in two treatments, (2) the difference in phenotypic value divided by the grand mean, and (3) the difference in phenotypic value divided by a pooled SD. Signal was present much more often when plasticity was measured as a simple difference (40% of comparisons) than when measured as a ratio (5% of comparisons in which the difference was divided by a grand mean or pooled SD). In this single case of a significant ratio ($[\text{Control} - \text{Crushed}/\text{SD}]$ for developmental stage), neither the numerator or denominator showed significant signal.

Studies of phylogenetic signal in plastic traits have been relatively rare. However there are studies comparing the plasticity of a few species (e.g., Cook 1968; Day et al. 1994; Smith and Van Buskirk 1995; Murren et al. 2015), studies containing a large number of species are less common, likely as a result of the chal-

lenge of conducting a large number of induction experiments. Of those studies that have been done, some have focused on testing the theoretical prediction that greater environmental heterogeneity is associated with greater amounts of plasticity by using phylogenetic contrasts (Van Buskirk 2002). Others have created phylogenies and mapped presence or absence of plasticity as a discrete trait onto the trees (Colbourne et al. 1997). In these cases, there do appear to be significant differences among major clades. Although these studies performed no explicit test for phylogenetic signal, it seems that phylogenetic signal likely existed. Examples include predator-induced morphological defenses in 34 species of zooplankton (*Daphnia*; Colbourne et al. 1997) and herbivore-induced defenses among 21 species of cotton (*Gossypium*; Thaler and Karban 1997). Finally, some studies have mapped different magnitudes of trait plasticity onto a phylogeny, but still not tested for significant phylogenetic signal. Most of these studies were focused on other research questions that simply needed phylogenetically corrected analyses. Examples include shade-induced and day-length-induced traits in six to nine species/populations of *Arabidopsis thaliana* (Pigliucci et al. 1999; Pollard et al. 2001) and temperature-induced changes in the larval period and morphology among 13 species of adult frogs (Gomez-Mestre and Buchholz 2006).

EVOLUTIONARY MODELS AND RATES OF TRAITS VERSUS TRAIT PLASTICITY

For every life-history trait, the OU model was either the best fitting model of character evolution or tied with BM based on AICc scores. The most common interpretation of the OU model is that it represents stabilizing selection due to the presence of adaptive optima (Butler et al. 2004, Beaulieu et al. 2012). The prominence of the OU model in our model fits may indicate that both traits and their plasticity tend to exhibit an adaptive optimum. We note, however, that in cases where α , the force of evolutionary attraction to an evolutionary optimum, is small, behavior of an OU model may be little different from BM (Cooper et al. 2016). Following the recommendation of Cooper et al. (2016), we used α to calculate “evolutionary half-life” (Hansen 1997) of each OU model in this study. This parameter does not have a strict biological interpretation, but Cooper et al. (2016) suggest that when it is large compared to the root age of a tree that it is estimated from, it reflects a mode of evolution that is more similar to BM than the OU model. Our results confirm that in cases where the OU model was a better fit than BM, that values of α were also large resulting in estimates of evolutionary half-life much less than the root age of the phylogeny that we used in our study (Table SA9). The EB model always showed the worst fit of the three models considered, indicating that evolutionary rates do not appear to be decreasing over time. This is in sharp contrast to what was found in studies of all mammals (Cooper and Davis 2010)

and birds (Brusatte et al. 2014), but similar to the results that Harmon et al. (2010) obtained when they examined smaller clades of animals.

We also examined the evolutionary rate of the traits versus the plasticity of the traits using ratio-transformed traits that yielded rate measures that generally differed by less than an order of magnitude. This is similar to the level of rate variation that Ackerly (2009) saw across plant traits (i.e., roughly two orders of magnitude) and that Rabosky and Adams (2012) observed among different clades for body mass. Based on these results, we suggest that ratio-transformed data, rather than raw or log-transformed data, should generally be used to compare the evolutionary rates of traits to trait plasticity, and when comparing rates among different measures of plasticity. Ratio transformation might also prove useful in general when calculating rates for traits that are not log-normal, or comparing the rates of lognormal and non-lognormal traits.

When we compared the evolutionary rate of traits versus trait plasticity using ratio transformation, the evolutionary rates of trait plasticity were greater than that of the traits from which they were calculated for three traits (time to hatching, developmental stage, and developmental rate) and slower for two life-history traits (mass at hatching and growth rate). This could be due to differences in the fitness consequence of phenotypic shifts in the direction or magnitude of plasticity compared to that of the traits themselves, or due to differences in the genetic architecture of traits and trait plasticity. Regardless, it seems that evolutionary responses of life-history traits to environmental changes or other selective pressures that push traits beyond or in opposition to their normal plasticity range could potentially happen more quickly through shifts in either trait means or through shifts in trait plasticity somewhat idiosyncratically depending upon the trait considered.

Conclusion

The results of this study demonstrate that plastic responses to environmental variation can be highly species-specific, which is not particularly surprising. More importantly, the traits that species possess consistently exhibit phylogenetic signal, yet the plasticity of those traits rarely exhibits phylogenetic signal. In addition, the OU model was either the best fitting model of character evolution or tied with BM for traits and all measures of trait plasticity, perhaps indicating the presence of adaptive peaks. Finally, whether trait plasticity evolves more slowly or faster than trait means varies depending upon the trait considered. Given the paucity of large studies that examine phylogenetic patterns in plasticity, it remains to be seen whether the patterns exhibited by amphibian embryos are representative of other taxa.

AUTHOR CONTRIBUTIONS

All authors except PRS executed the experiments. RAR, PRS, and JIH analyzed the data and wrote the paper. RAR and PRS secured funding for the project.

ACKNOWLEDGMENTS

This project was funded by a National Science Foundation grant (DEB 07-16149) to RAR and PRS. During the data analysis, JIH was supported by National Institute of General Medical Sciences Grant K12GM088021. Content is solely the responsibility of the authors and does not necessarily represent the official views of either the NIH or NSF. We were also assisted by a large number of undergraduate students spanning all of the laboratories. The authors declare no conflicts of interest. We also thank the Associate Editor and three excellent reviewers for their help in making this a stronger, clearer, and more compelling publication.

DATA ARCHIVING

The doi for our data is <https://doi.org/10.5061/dryad.7p02c>.

LITERATURE CITED

- Ackerly, D. 2009. Conservatism and diversification of plant functional traits: evolutionary rates versus phylogenetic signal. *Proc. Natl. Acad. Sci. USA* 106:19699–19706.
- Anderson, A. L., and W. D. Brown. 2009. Plasticity of hatching in green frogs (*Rana clamitans*) to both egg and tadpole predators. *Herpetologica* 65:207–213.
- Anderson A. R., and J. W. Petranka. 2003. Odonate predator does not affect hatching time or morphology of two amphibians. *J. Herpetol.* 37:65–71.
- Beaulieu, J. M., D. C. Jhwueng, C. Boettiger, and B. C. O'Meara. 2012. Modeling stabilizing selection: expanding the Ornstein–Uhlenbeck model of adaptive evolution. *Evolution* 66:2369–2383.
- Berendonk, T. U., T. G. Barraclough, and J. C. Barraclough. 2003. Phylogenetics of pond and lake lifestyles in *Chaoborus* midge larvae. *Evolution* 57:2173–2178.
- Blomberg, S. P., T. Garland Jr., and A. R. Ives. 2003. Testing for phylogenetic signal in comparative data: behavioral traits are more labile. *Evolution* 57:717–745.
- Blomberg, S. P., and T. Garland Jr., 2002. Tempo and mode in evolution: Phylogenetic inertia, adaptation and comparative methods. *J. Evol. Biol.* 15:899–910.
- Boersma, M., P. Spaak, and L. De Meester. 1998. Predator-mediated plasticity in morphology, life history, and behavior of *Daphnia*: uncoupling of responses. *Am. Nat.* 152:237–248.
- Brandt, R., and C. A. Navas. 2011. Life history evolution on tropidurinae lizards: influence of lineage, climate, and body size. *PLoS One* 6:e20040.
- Brusatte, S.L., G. T. Lloyd, S. C. Wang, and M. A. Norell. 2014. Gradual assembly of avian body plan culminated in rapid rates of evolution across the dinosaur-bird transition. *Curr. Biol.* 24:2386–2392.
- Burns, J. H., and S. Y. Strauss. 2012. Effects of competition on phylogenetic signal and phenotypic plasticity in plant functional traits. *Ecology* 93:S126–S137.
- Butler, M. A., and A. A. King. 2004. Phylogenetic comparative analysis: a modeling approach for adaptive evolution. *Am. Nat.* 164:683–695.
- Cadle, J. E., and H. W. Greene. 1993. Phylogenetic patterns, biogeography, and the ecological structure of neotropical snake assemblages. Pp. 281–293 in R. E. Ricklefs and D. Schluter, eds. *Species diversity in ecological communities: historical and geographical perspectives*. Univ. of Chicago Press, Chicago, IL.

- Callahan, H. S., H. Maughan, and U. K. Steiner. 2008. Phenotypic plasticity, costs of phenotypes, and costs of plasticity. *Ann. N Y Acad. Sci.* 1133:44–66.
- Chivers, D. P., J. M. Kiesecker, A. Marco, J. DeVito, M. T. Anderson, and A. R. Blaustein. 2001. Predator-induced life history changes in amphibians: egg predation induces hatching. *Oikos* 92:135–142.
- Colbourne, J. K., P. D. N. Hebert, and D. J. Taylor. 1997. Evolutionary origins of phenotypic diversity in *Daphnia*. Pp. 163–188 in T. J. Givnish and K. J. Sytsma, eds. *Molecular evolution and adaptive radiation*. Cambridge Univ. Press, Cambridge, U.K.
- Cook, C. D. K. 1968. Phenotypic plasticity with particular reference to three amphibious plant species. Pp. 97–111 in V. H. Heywood, ed. *Modern methods in plant taxonomy*. Academic Press, London.
- Cooper, N., and A. Purvis. 2010. Body size evolution in mammals: complexity in tempo and mode. *Am. Nat.* 175:727–738.
- Cooper, N., G. H. Thomas, and R. G. FitzJohn. 2016. Shedding light on the “dark side” of phylogenetic comparative methods. *Methods Ecol. Evol.* 7:693–699.
- Day, T., J. Pritchard, and D. Schluter. 1994. A comparison of two sticklebacks. *Evolution* 48:1723–1734.
- DeWitt, T. J., and S. M. Scheiner. 2004. *Phenotypic plasticity: functional and conceptual approaches*. Oxford Univ. Press, New York.
- Dibble, C. J., J. E. Kauffman, E. M. Zuzik, G. R. Smith, and J. E. Rettig. 2009. Effects of potential predator and competitor cues and sibship on wood frog (*Rana sylvatica*) embryos. *Amphibia Reptilia* 30:294–298.
- Diggle, P. K. 2002. A developmental morphologist’s perspective on plasticity. *Evol. Ecol.* 16:267–283.
- Doughty, P. 1995. Testing the ecological correlates of phenotypically plastic traits within a phylogenetic framework. *Acta Oecol.* 16:519–524.
- Felsenstein, J. 2004. *Inferring phylogenies*. Sinauer Associates, Sutherland, MA.
- Gomez-Mestre, I., and D. Buchholz. 2006. Developmental plasticity mirrors differences among taxa in spadefoot toads linking plasticity and diversity. *Proc. Natl. Acad. Sci. USA* 103:19021–19026.
- Gomez-Mestre, I., J. J. Wiens, and K. M. Warkentin. 2008. Evolution of adaptive plasticity: risk-sensitive hatching in Neotropical leaf-breeding treefrogs. *Ecol. Monogr.* 78:205–224.
- Gosner, K. L. 1960. A simplified table for staging anuran embryos and larvae with notes on identification. *Herpetologica* 16:183–190.
- Grant, P. R. 1986. *Ecology and evolution of Darwin’s finches*. Princeton Univ. Press, Princeton, NJ.
- Harmon, L. J., J. T. Weir, C. D. Brock, R. E. Glor, and W. Challenger. 2008. GEIGER: investigating evolutionary radiations. *Bioinformatics* 24:129–131.
- Harmon, L. J., J. B. Losos, T. J. Davies, G. C. Gillespie, J. L. Gittleman, W. B. Jennings, K. H. Kozak, M. A. McPeck, F. Moreno-Roark, T. J. Near, et al. 2010. Early bursts of body size and shape evolution are rare in comparative data. *Evolution* 64:2385–2396.
- Harvey, P. H., and M. D. Pagel. 1991. *The comparative method in evolutionary biology*. Oxford Univ. Press, New York.
- Hedges, L. V. 1981. Distribution theory for Glass’ estimator of effect size and related estimators. *J. Educ. Stat.* 6:107–128.
- Johnson, J. B., D. Saenz, C. K. Adams, and R. N. Conner. 2003. The influence of predator threat on the timing of a life history switch point: predator-induced hatching in the southern leopard frog (*Rana sphenoccephala*). *Can. J. Zool.* 81:1608–1613.
- Kembel, S. W., P. D. Cowan, M. R. Helmus, W. K. Cornwell, H. Morlon, D. D. Ackerly, S. P. Blomberg, and C. O. Webb. 2010. Picante: R tools for integrating phylogenies and ecology. *Bioinformatics* 26:1463–1464.
- Laurila, A., P.-A. Crochet, and J. Merilä. 2001. Predation-induced effects on hatching morphology in the common frog (*Rana temporaria*). *Can. J. Zool.* 79:926–930.
- Laurila, A., S. Pakkasmaa, P. A. Crochet, and J. Merila. 2002. Predator-induced plasticity in early life history and morphology in two anuran amphibians. *Oecologia* 132:524–530.
- Losos, J. B. 1990. A phylogenetic analysis of character displacement in Caribbean *Anolis* communities. *Evolution* 44:558–569.
- . 2008. Phylogenetic niche conservatism, phylogenetic signal and the relationship between phylogenetic relatedness and ecological similarity among species. *Ecol. Lett.* 11:995–1003.
- Losos, J. B., T. R. Jackman, A. Larson, K. de Queiroz, and L. Rodriguez-Schettino. 1998. Contingency and determinism in replicated adaptive radiations of island lizards. *Science* 279:2115–2118.
- Mandrillon, A.-L., and P. Saglio. 2007. Effects of embryonic exposure to conspecific chemical cues on hatching and larval traits in the common frog (*Rana temporaria*). *Chemoecology* 17:169–175.
- McPeck, M. A., and J. M. Brown. 2000. Building a regional species pool: Diversification of the *Enallagma* damselflies in eastern North American waters. *Ecology* 81:904–920.
- Miner, B. G., S. E. Sultan, S. G. Morgan, D. K. Padilla, and R. A. Relyea. 2005. Ecological consequences of phenotypic plasticity. *Trends Ecol. Evol.* 20:685–692.
- Moles, A. T., D. D. Ackerly, C. O. Webb, J. C. Tweddle, J. B. Dickie, and M. Westoby. 2005. A brief history of seed size. *Science* 307:576–580.
- Murren, C. J., H. J. Maclean, S. E. Diamond, U. K. Steiner, M. A. Heskell, C. A. Handelsman, C. K. Ghalambor, J. R. Auld, H. S. Callahan, D. W. Pfennig, et al. 2014. Evolutionary change in continuous reaction norms. *Am. Nat.* 183:453–467.
- Murren, C. J., J. R. Auld, H. Callahan, C. K. Ghalambor, C. A. Handelsman, M. A. Heskell, J. G. Kingsolver, H. J. Maclean, J. Masel, H. Maughan, et al. 2015. Constraints on the evolution of phenotypic plasticity: costs of phenotype and costs of plasticity. *Heredity* 115:293–301.
- Orizaola, G., and F. Braña. 2004. Hatching responses of four newt species to predatory fish chemical cues. *Ann. Zool. Fennici* 41:635–645.
- Pennell, M. W., J. M. Eastman, G. J. Slater, J. W. Brown, J. C. Uyeda, R. G. FitzJohn, M. E. Alfaro, and L. J. Harmon. 2014. Geiger v2. 0: an expanded suite of methods for fitting macroevolutionary models to phylogenetic trees. *Bioinformatics* 30:2216–2218.
- Pigliucci M. 2001. Phenotypic integrations: studying the ecology and evolution of complex phenotypes. *Ecol. Lett.* 6:256–272.
- Pigliucci, M., K. Cammell, and J. Schmitt. 1999. Evolution of phenotypic plasticity a comparative approach in the phylogenetic neighbourhood of *Arabidopsis thaliana*. *J. Evol. Biol.* 12:779–791.
- Pigliucci, M., H. Pollard, and M. B. Cruzan. 2003. Comparative studies of evolutionary responses to light environments in *Arabidopsis*. *Am. Nat.* 161:68–82.
- Pollard, H., M. Cruzan, and M. Pigliucci. 2001. Comparative studies of reaction norms in *Arabidopsis*. I. Evolution of responses to day length. *Evol. Ecol. Res.* 3:129–155.
- Poo, S., and D. P. Bickford. 2014. Hatching plasticity in a Southeast Asian tree frog. *Behav. Ecol. Sociobiol.* 68:1733–1740.
- Price, T., I. J. Lovette, E. Bermingham, H. L. Gibbs, and A. D. Richman. 2000. The imprint of history on communities of North American and Asian warblers. *Am. Nat.* 156:354–367.
- Pyron, R. A., and J. J. Wiens. 2013. Large-scale phylogenetic analyses reveal the causes of high tropical amphibian diversity. *Proc. R. Soc. Lond.* 280. <https://doi.org/10.1098/rspb.2013.1622>.

- Rabosky, D. L., and D. C. Adams. 2012. Rates of morphological evolution are correlated with species richness in salamanders. *Evolution* 66:1807–1818.
- Relyea, R. A. 2007. Getting out alive: how predators affect the decision to metamorphose. *Oecologia* 152:389–400.
- Revell, L. J., L. J. Harmon, and D. C. Collar. 2008. Phylogenetic signal, evolutionary process, and rate. *Syst. Biol.* 57:591–601.
- Richardson, J. M. L. 2001a. A comparative study of activity levels in larval anurans and response to the presence of different predators. *Behav. Ecol.* 12:51–58.
- . 2001b. The relative roles of adaptation and phylogeny in determination of larval traits in diversifying anuran lineages. *Am. Nat.* 157:282–299.
- . 2002a. Burst swim speed in tadpoles inhabiting ponds with different top predators. *Evol. Ecol. Res.* 4:627–642.
- . 2002b. A comparative study of phenotypic traits related to resource utilization in anuran communities. *Evol. Ecol.* 16:101–122.
- Saenz, D., J. B. Johnson, C. K. Adams, and G. H. Dayton. 2003. Accelerated hatching of southern leopard frog (*Rana sphenoccephala*) eggs in response to the presence of a crayfish (*Procambarus nigrocinctus*) predator. *Copeia* 2003:646–649.
- Schalk, G., M. R. Forbes, and P. J. Weatherhead. 2002. Developmental plasticity and growth rates of green frog (*Rana clamitans*) embryos and tadpoles in relation to a leech (*Macrobdella decora*) predator. *Copeia* 2002:445–449.
- Schlichting, C., and M. Pigliucci. 1998. Phenotypic evolution: a reaction norm perspective. Sinauer, Sunderland.
- Schluter, D. 2000. The ecology of adaptive radiation. Oxford Univ. Press, New York.
- Schoeppner, N. M., and R. A. Relyea. 2008. Detecting small environmental differences: Risk-response curves for predator-induced behavior and morphology. *Oecologia* 154:743–754.
- Segev, O., A. Rodríguez, S. Hauswaldt, K. Hugemann, and M. Vences. 2015. Flatworms (*Schmidtea nova*) prey upon embryos of the common frog (*Rana temporaria*) and induce minor developmental acceleration. *Amphibia Reptilia* 36:155–163.
- Smith, D. C., and J. Van Buskirk. 1995. Phenotypic design, plasticity, and ecological performance in two tadpole species. *Am. Nat.* 145:211–233.
- Sota, T., and R. Ishikawa. 2004. Phylogeny and life-history evolution in *Carabus* (subtribe Carabina: Coleoptera, Carabidae) based on sequences of two nuclear genes. *Biol. J. Linnean Soc.* 81:135–149.
- Special Feature of Ecology. 2012. <http://esajournals.onlinelibrary.wiley.com/hub/issue/10.1002/ecy.2012.93.issue-sp8/>.
- Stephens, P. R., and J. J. Wiens. 2004. Convergence, divergence, and homogenization in the ecological structure of emydid turtle communities: the effects of phylogeny and dispersal. *Am. Nat.* 164:244–254.
- Thaler, J. S., and R. Karban. 1997. A phylogenetic reconstruction of constitutive and induced resistance in *Gossypium*. *Am. Nat.* 149:1139–1146.
- Touchon, J. C., I. Gomez-Mestre, and K. M. Warkentin. Hatching plasticity in tow temperate anurans: response to a pathogen and predation cues. *Can. J. Zool.* 84:556–563.
- Van Buskirk, J. 2002. A comparative test of the adaptive plasticity hypothesis: relationships between habitat and phenotype in anuran larvae. *Am. Nat.* 160:87–102.
- Vonesh, J. R. 2005. Egg predation and predator-induced hatching plasticity in the African reed frog, *Hyperolius spinigularis*. *Oikos* 110:241–252.
- Warkentin, K. M. 1995. Adaptive plasticity in hatching age: a response to predation risk trade-offs. *Proc. Natl. Acad. Sci. USA* 92:3507–3510.
- Webb, C. O., D. D. Ackerly, M. A. McPeck, and M. J. Donoghue. 2002. Phylogenies and community ecology. *Annu. Rev. Ecol. Syst.* 33:475–505.
- West-Eberhard, M. J. 2003. Developmental plasticity and evolution. Oxford Univ. Press, New York.
- Winemiller, K. O. 1991. Ecomorphological diversification in lowland freshwater fish assemblages from five biotic regions. *Ecol. Monogr.* 61:343–365.

Associate Editor: D. Rabosky
Handling Editor: M. Noor

Supporting Information

Additional Supporting Information may be found in the online version of this article at the publisher's website:

Table SA1. Locations of amphibians collected for the embryonic plasticity experiments, the number of clutches collected, and the developmental stage at which each experiment began.

Table SA2. MANOVA results from the analysis of traits measured in the embryonic experiments across 23 species of amphibians.

Table SA3. MANOVA results from the analysis of traits measured in the embryonic experiments across three species of amphibians conducted at two temperatures.

Table SA4. Tests of evolutionary rates (σ^2) for time to hatching of amphibian embryos when raised in three inducing environments.

Table SA5. Tests of evolutionary rates (σ^2) for mass at hatching of amphibian embryos when raised in three inducing environments.

Table SA6. Tests of evolutionary rates (σ^2) for developmental stage at hatching of amphibian embryos when raised in three inducing environments.

Table SA7. Tests of evolutionary rates (σ^2) for growth rate of amphibian embryos when raised in three inducing environments.

Table SA8. Tests of evolutionary rates (σ^2) for developmental rate of amphibian embryos when raised in three inducing environments.

Figure SA1. Chronogram used for phylogenetic analyses obtained from Pyron and Wiens (2013).

Figure SA2. A comparison of evolutionary rate versus trait range for raw values of developmental rate.